arXiv:hep-th/9212092v1 15 Dec 1992

Self-Duality and Supersymmetry
E.Sezgin
USE PLAIN TEX

CTP-TAMU-85/92
hepth@xxx/9212092
December 1992

Self-Duality and Supersymmetry

E. Sezgin *

Center for Theoretical Physics, Texas A&M University,
College Station, Texas 77843-4242, USA.

ABSTRACT

Chiral/self-dual restrictions of various super Yang-Mills and supergravity theories in (2,2) dimensions are described. These include the N=1 supergravity with a cosmological term and the N=1 new minimal supergravity theory. In the latter case, a self-duality condition on a torsionful Riemann curvature is possible, and it implies the equations of motion that follow from an R^2 type supergravity action.

---

* Supported in part by NSF, under grant PHY-9106593.
Contribution to the Proceedings of the Trieste Summer Workshop on Superstrings and Related Topics, Trieste, July 1992.

1. Self-dual Yang-Mills and self-dual gravity equations in (2,2) dimensions are known to arise in open and closed N=2 string theories [1], i.e. string theories with *local* N=2 worldsheet supersymmetry. Recently, it has been shown that N=2 open string actually describes a self-dual *super* Yang-Mills theory, while the N=2 closed string describes self-dual N=8 supergravity theory and the heterotic N=2 closed string describes *gauged* N=8 supergravity theory in (2,2) dimensions [2]. These interesting developments motivate further study of various self-dual supersymmetric field theories in (2,2) dimensions.

Another motivation for studying the self-dual systems in (2,2) dimensions is their connection with a large class of integrable systems in (1,1) dimensions [3]. For example, different types of dimensional reductions of the (2,2) dimensional self-dual Yang-Mills theory yield a large class of integrable systems in (1,1) dimensions including the KdV equation, the non-linear Schrodinger equations and Toda equations. One expects that even a larger class of integrable models are e mbedded in self-dual supersymmetric field theories. In fact, it has recently been shown that super KdV and super-Toda equations do arise from dimensional reductions of N=2 supersymmetric self-dual Yang-Mills system in (2,2) dimensions [4].

In this note, we shall describe the algebraic structure of various self-dual supersymmetric field theories in (2,2) dimensions. We first describe their underlying superalgebras. Next, we discuss the chiral/self-dual restrictions of N=1,2,4 super Yang-Mills theories. Finally, we describe the chiral/self-dual restrictions of (i) N=1 supergravity (ii) new minimal N=1 supergravity and (iii) N=1 supergravity with a cosmological constant in (2,2) dimensions. In the second case, we find a self-duality condition on a torsionful Riemann curvature which implies the equations of motion that follow from an R^2 type action [5]. In the last case, surprisingly, it turns out that (a) the self-duality conditions on the Riemann tensor and the gravitino curvature are precisely those of self-dual N=1 supergravity without a cosmological constant, and (b) in the chiral truncation of the supersymmetry transformation rules of anti de Sitter supergravity, the cosmological constant parameter survives. Thus, the usual self-dual supergravity equations seem to be invariant under a one parameter extension of the super-Poincaré group which is a particular contraction of the anti de Sitter Poincaré algebra in (2,2) dimensions. A similar phenomenon has been descibred in the context of gauged N=8 supergravity in [2].

2. Let us consider first the relevant super Poincaré algebras. A characteristic feature of the super Poincaré algebra in (2,2) dimensions is that the supercharges are independent real Majorana-Weyl spinors [6]. Various supermultiplets in (3,1) dimensions can easily be Wick rotated to supermultiplets in (2,2) dimensions. Of particular interest is the Yang-Mills supermultiplet consisting of the Yang-Mills gauge field and its fermionic partner. We recall that this supermultiplet also exists in (5,1), where the supercharges are pseudo-real Majorana-Weyl, and in (9,1) dimensions where the supercharges are real Majorana-Weyl. We can Wick rotate the first one to (3,3) dimensions, and the latter to (5,5) dimensions. In both cases, there exist real Majorana-Weyl spinors. Thus, it is useful to consider the (2,2), (3,3), (5,5) and (9,1) dimensions, where real Majorana-Weyl spinors and simple Yang-Mills supermultiplets exist.

In (2,2) dimensions, we adopt the following conventions. The spacetime signature is $(--++)$, the charge conjugation matrix is $C=\gamma_{12}$, $C^T=-C$, $(\gamma_{\mu\nu}C)^T=\gamma_{\mu\nu}C$, $\gamma_5=\gamma_{1234}$, $\gamma_5^T=\gamma_5$, $[C,\gamma_5]=0$, $\gamma_\mu^*=\eta\gamma_\mu$, and $(\gamma_\mu C)^T=-\eta\gamma_\mu C$, where $\eta=\pm1$. Without loss of generality, we shall set $\eta=1$. The extended Poincaré algebra consists of Poincaré generators and supercharges $Q_\pm^i$, $i=1,...,N$ which obey the anticommutation rules

$$\begin{aligned}
&\{Q_+^i,Q_-^j\}=\tfrac{1}{2}(1+\gamma_5)\gamma^\mu CP_\mu\lambda^{ij}\ ,\\
&\{Q_\pm^i,Q_\pm^j\}=0\ ,
\end{aligned}\tag{1}$$

where λ_{ij} can be chosen to be either the $SO(n,N-n)$ invariant symmetric tensor η_{ij} or the $Sp(n,N-n)$ invariant antisymmetric tensor Ω_{ij}. These are the automorphism groups of the algebra, and the supercharges transform in their defining representations.

In (3,3) dimensions, we adopt the following conventions: The spacetime signature is $(---+++)$, the charge conjugation matrix is $C=\gamma_{123}$, $(\gamma_\mu C)^T=-(\gamma_\mu C)$, $(\gamma_{\mu\nu\rho}C)^T=\gamma_{\mu\nu\rho}C$, $\gamma_7=\gamma_{123456}$, $\gamma_7^T=\gamma_7$, $\{C,\gamma_7\}=0$, $\gamma_\mu^*=\eta\gamma_\mu$, $C^T=-\eta C$ and $(\gamma_{\mu\nu}C)^T=-\eta\gamma_{\mu\nu}C$, where $\eta=\pm1$. Without loss of generality, we shall set $\eta=1$. The supercharges Q_+^i , $i=1,...,M$ and Q_-^a, $a=1,...,N$ obey the anticommutation rules

$$\begin{aligned}
&\{Q_+^i,Q_+^j\}=\tfrac{1}{2}(1+\gamma_7)\gamma^\mu CP_\mu\Omega^{ij}\ ,\\
&\{Q_+^i,Q_-^a\}=0\ ,\\
&\{Q_+^a,Q_+^b\}=\tfrac{1}{2}(1+\gamma_7)\gamma^\mu CP_\mu\Omega^{ab}\ ,
\end{aligned}\tag{2}$$

where Ω^{ij} and Ω^{ab} are the invariant tensors of the automorphism group $Sp(m,M-m)\times Sp(n,N-n)$.

Finally, in (5,5) dimensions, we adopt the following conventions. The spacetime signature is $(-----+++++)$, the charge conjugation matrix is $C=\gamma_{12345}$, $(\gamma_\mu C)^T=(\gamma_\mu C)$, $(\gamma_{\mu\nu\rho}C)^T=-\gamma_{\mu\nu\rho}C$, $(\gamma_{\mu\nu\rho\lambda\tau}C)^T=\gamma_{\mu\nu\rho\lambda\tau}C$, $\gamma_{11}=\gamma_{1\ldots10}$, $\gamma_{11}^2=1$, $\gamma_{11}^T=\gamma_{11}$, $\gamma_{11}^*=\gamma_{11}$, $\{C,\gamma_{11}\}=0$, $\gamma_\mu^*=\eta\gamma_\mu$, $C^T=-\eta C$, $(\gamma_{\mu\nu}C)^T=\eta\gamma_{\mu\nu}C$, and $(\gamma_{\mu\nu\rho\sigma}C)^T=-\eta\gamma_{\mu\nu\rho\sigma}C$, where $\eta=\pm1$. Again, without loss of generality, we shall set $\eta=1$. The supercharges Q_+^i , $i=1,...,M$ and Q_-^a, $a=1,...,N$ obey the anticommutation rules

$$\begin{aligned}
&\{Q_+^i,Q_+^j\}=\tfrac{1}{2}(1+\gamma_{11})\gamma^\mu CP_\mu\eta^{ij}\ ,\\
&\{Q_+^i,Q_-^a\}=0\ ,\\
&\{Q_+^a,Q_+^b\}=\tfrac{1}{2}(1+\gamma_{11})\gamma^\mu CP_\mu\eta^{ab}\ ,
\end{aligned}\tag{3}$$

where η^{ij} and η^{ab} are the invariant tensors of the automorphism group $SO(m,M-m)\times SO(n,N-n)$. The properties of the Dirac matrices and the form of the extended Poincaré algebra in $(9,1)$ dimensions is the same as in the $(5,5)$ dimensions described above.

3. We now turn to the description of self-dual super Yang-Mills theories in $(2,2)$ dimensions. It can easily obtained by a Wick rotation of the super Yang-Mills theory in $(3,1)$ dimensions. The model contains the gauge field A_μ and its fermionic parners $\lambda_\pm$, which are now independent real Majorana-Weyl spinors. The self-dual version of the theory is obtained by setting $\lambda_+ = 0$ [7]. The resulting transformation rules are

$$\begin{aligned} \delta A_\mu &= \tfrac{1}{2}\bar\epsilon_+\gamma_\mu\lambda_- \ , \\ \delta\lambda_- &= -\tfrac{1}{4}\gamma^{\mu\nu}F_{\mu\nu}\epsilon_- \ . \end{aligned} \tag{4}$$

Note that both supersymmetry parameters $\epsilon_\pm$ are present. The closure of these transformations then requires the conditions

$$\begin{aligned} F_{\mu\nu} - \tfrac{1}{2}\epsilon_{\mu\nu\rho\sigma}F^{\rho\sigma} &= 0, \\ \gamma^\mu D_\mu\lambda_- &= 0 \ , \end{aligned} \tag{5}$$

where D_μ is the gauge covariant derivative. These equations transform into each other under the supersymmetry transformations: $\delta F^-_{\mu\nu} = -\frac{1}{2}\bar\epsilon_+\gamma_{\mu\nu}\gamma^\rho D_\rho\lambda_-$ and $\delta\gamma^\mu D_\mu\lambda_- = -\frac{1}{4}\gamma^\mu\gamma^{\nu\rho}D_\mu F^-_{\nu\rho}\epsilon_-$, where $F^-_{\mu\nu} \equiv F_{\mu\nu} - \frac{1}{2}\epsilon_{\mu\nu\rho\sigma}F^{\rho\sigma}$.

As to be expected, the N=1 self-dual super Yang-Mills equations (5) are also invariant the transformations of the superconformal group $SL(4|1)$ whose bosonic subgroup is $SL(4,R)(\approx SO(3,3)) \times SO(1,1)$. The superconformal transformations are given by (4) and

$$\begin{aligned} \delta A_\mu &= -\tfrac{1}{2}\bar\eta_-\gamma^\nu\gamma_\mu\lambda_- x_\nu + \xi^\lambda\partial_\lambda A_\mu \ , \\ \delta\lambda_- &= -\tfrac{1}{4}\gamma^{\mu\nu}\gamma^\rho x_\rho F_{\mu\nu}\eta_+ + \tfrac{3}{2}\alpha\lambda_- - \beta\lambda_- + \tfrac{1}{4}\omega^{\mu\nu}\gamma_{\mu\nu}\lambda_- + \xi^\mu\partial_\mu\lambda_- \ , \end{aligned} \tag{6}$$

where $\eta_\pm$ are the special conformal supersymmetry parameters, α, β are the parameters of dilatations and SO(1,1) transformations, respectively, and ξ^μ represent the spacetime conformal transformations given by

$$\xi^\mu(x) = a^\mu - \omega^{\mu\nu}x_\nu + \alpha x^\mu + 2x^\mu\eta\cdot x - \eta^\mu x^2 \ . \tag{7}$$

Here $a^\mu, \omega^{\mu\nu}$ and η^μ are the constant parameters of translations, Lorentz rotations and conformal boosts, respectively.

4. The on-shell N=2 super Yang-Mills multiplet in (2,2) dimensions contains the fields $(A_\mu, \lambda^i_\pm, S, \phi)$ where $i = 1,2$ and S, ϕ are two real scalars, all taking their values in the adjoint representation of some Lie algebra. It is easiest to obtain the transformation rules for this multiplet from (3,3) dimensions by ordinary dimensional reduction. The simple super Yang-Mills multiplet transformation rules in (3,3) dimensions are

$$\begin{aligned} \delta A_{\hat\mu} &= \tfrac{1}{2}\bar\eta^i\Gamma_{\hat\mu}\chi^j\epsilon_{ij} \ , \qquad\qquad \hat\mu = 1, ..., 6 \\ \delta\chi^i &= -\tfrac{1}{4}\Gamma^{\hat\mu\hat\nu}F_{\hat\mu\hat\nu}\eta^i \ , \end{aligned} \tag{8}$$

where $\Gamma_7\chi = -\chi$ and $\Gamma_7\eta = -\eta$. Let us choose the Dirac matrices as $\Gamma^\mu = \gamma^\mu \times 1$, $\mu = 1,2,5,6$, $\Gamma_3 = \gamma_5 \times i\sigma_2$, $\Gamma_4 = \gamma_5 \times \sigma_1$. Performing a straigtforward dimensional reduction in which we take all the fields to be independent of the two internal coordinates and defining $\phi \equiv \frac{1}{2}(A_3 - A_4)$, and $S \equiv \frac{1}{2}(A_3 + A_4)$, from (8) we find the transformation rules for the $N = 2$ super Yang-Mills multiplet in (2,2) dimensions:

$$\begin{aligned}
\delta A_\mu &= \tfrac{1}{2}\bar{\eta}^i_+ \gamma_\mu i\sigma_2 \chi^j_- \epsilon_{ij} - \tfrac{1}{2}\bar{\eta}^i_- \gamma_\mu i\sigma_2 \chi^j_+ \epsilon_{ij} \ , \\
\delta\phi &= -\tfrac{1}{2}\bar{\eta}^i_- \chi^j_- \epsilon_{ij} \ , \\
\delta S &= -\tfrac{1}{2}\bar{\eta}^i_+ \chi^j_+ \epsilon_{ij} \ , \\
\delta\chi^i_+ &= -\tfrac{1}{4}\gamma^{\mu\nu} F_{\mu\nu}\eta^i_+ - \gamma^\mu D_\mu\phi\sigma_+\eta^i_- + \gamma^\mu D_\mu S\sigma_-\bar{\eta}^i_- - [S,\phi]\eta^i_+ \ , \\
\delta\chi^i_- &= -\tfrac{1}{4}\gamma^{\mu\nu} F_{\mu\nu}\eta^i_- + \gamma^\mu D_\mu\phi\sigma_+\eta^i_+ - \gamma^\mu D_\mu S\sigma_-\bar{\eta}^i_- + [S,\phi]\eta^i_+ \ ,
\end{aligned} \tag{9}$$

where $\gamma_5\chi^i_\pm = \pm\chi^i_\pm$ and $\sigma_\pm = \frac{1}{2}(\sigma_1 \pm \mathrm{i}\sigma_2)$. To obtain the self-dual version, we impose the conditions [8]

$$S = 0 \ , \qquad \chi^i_+ = 0 \ , \qquad \sigma_+\chi^i_- = 0 \ , \qquad \sigma_+\eta^i_- = 0 \ , \qquad \sigma_-\eta^i_+ = 0 \ . \tag{10}$$

Using the notation, $\sigma_-\chi^i_- \equiv \lambda^i_-$ and $\sigma_+\eta^i_+ \equiv \epsilon^i_+$, we then have

$$\begin{aligned}
\delta A_\mu &= \tfrac{1}{2}\bar{\epsilon}^i_+ \gamma_\mu \lambda^j_- \epsilon_{ij} \ , \\
\delta\phi &= -\tfrac{1}{2}\bar{\epsilon}^i_- \lambda^j_- \epsilon_{ij} \ , \\
\delta\lambda^i_- &= -\tfrac{1}{4}\gamma^{\mu\nu} F_{\mu\nu}\epsilon^i_- + \gamma^\mu D_\mu\phi\epsilon^i_+ \ .
\end{aligned} \tag{11}$$

The closure of the algebra modulo a set of equations of motions that transform into each other under superymmetry requires the following set of equations [8]

$$\begin{aligned}
F_{\mu\nu} - \tfrac{1}{2}\epsilon_{\mu\nu\rho\sigma}F^{\rho\sigma} &= 0 \ , \\
\gamma^\mu D_\mu \lambda^i_- &= 0, \\
D_\mu D^\mu \phi - \tfrac{1}{8}\epsilon_{ij}[\bar{\lambda}^i_-, \lambda^j_-] &= 0 \ ,
\end{aligned} \tag{12}$$

where the commutator is in the space of the Lie algebra generators. The last equation is easily seen to follow from the supersymmetric variation of the second equation. One application of these equations has already been considered in [4] where it has been an appropriate dimensional reduction of these equations to (1,1) dimensions yields the super KdV equations.

5. We now consider the case of N=4 super Yang-Mills theory. While it can be formulated in (2,2) dimensions, it does not admit a simple self-dual truncation of the kind discussed above. The N=4 super Yang-Mills theory in (2,2) dimensions can be easily obtained by an ordinary dimensional reduction from the simple super Yang-Mills theory in (5,5) dimensions, whose transformation rules are

$$\begin{aligned} \delta A_{\hat\mu} &= \tfrac{1}{2}\bar\epsilon\Gamma_{\hat\mu}\lambda\ , \qquad\qquad \hat\mu = 1, ..., 10 \\ \delta\lambda &= -\tfrac{1}{4}\Gamma^{\hat\mu\hat\nu}F_{\hat\mu\hat\nu}\epsilon\ , \end{aligned} \tag{13}$$

where $\Gamma_{11}\lambda = -\lambda$ and $\Gamma_{11}\epsilon = -\epsilon$. Let us choose the Dirac matrices as $\Gamma_\mu = \gamma_\mu \times 1$ $(\mu = 1, 2, 9, 10)$ and $\Gamma_i = \gamma_5 \times \gamma_i$ $(i = 3, ..., 8)$. As usual, setting $\partial_i = 0$, we obtain the N=4 super Yang-Mills transformation rules

$$\begin{aligned} \delta A_\mu &= \bar\epsilon_+\gamma_\mu\lambda_- + \bar\epsilon_-\gamma_\mu\lambda_+\ , \\ \delta\phi_i &= -\bar\epsilon_-\gamma_i\lambda_- + \bar\epsilon_+\gamma_i\lambda_+\ , \\ \delta\lambda_+ &= -\tfrac{1}{4}F^{\mu\nu}\gamma_{\mu\nu}\epsilon_+ - \tfrac{1}{2}\gamma^\mu\gamma^i D_\mu\phi_i\epsilon_- - \tfrac{1}{4}\gamma^{ij}[\phi_i, \phi_j]\epsilon_+\ , \\ \delta\lambda_- &= -\tfrac{1}{4}F^{\mu\nu}\gamma_{\mu\nu}\epsilon_- + \tfrac{1}{2}\gamma^\mu\gamma^i D_\mu\phi_i\epsilon_+ - \tfrac{1}{4}\gamma^{ij}[\phi_i, \phi_j]\epsilon_-\ , \end{aligned} \tag{14}$$

where ϵ and λ carry the spinor indices of the Lorentz group SO(2,2) as well as the spinor indices of the internal symetry group SO(3,3), $\bar\epsilon = \epsilon^T C_4 C_6$ with $C_4 = \gamma_{12}$, $C_6 = \gamma_{345}$, $A_i \equiv \phi_i$, $\gamma_5\epsilon_\pm = \pm\epsilon_\pm$ and $\gamma_7\epsilon_\pm = \mp\epsilon_\pm$. Now it can be seen that λ_+ can not be consistently set equal to zero in order to arrive at the self-duality equation and its super-partners. Unlike in the case of N=2 super Yang-Mills, here the second and third terms in the transformation rule of λ_+ can not be made to vanish by any condition on the scalar fields or the parameters $\epsilon_\pm$ in a way consistent with supersymmetry [2,8]. A different kind of self-dual N=4 super Yang-Mills theory has been formulated in [2], where it is also shown that it arises as the effective theory in the open N=2 superstring theory. In this version, one introduces an anti-self-dual Lagrange multiplier field whose equation of motion imposes the self-duality condition on the Yang-Mills curvature, and an anti-chiral spinor. The Lagrange multiplier field propagates, however, and consequently the number of helicities in the theory do not get halved.

6. We now turn to the description of two types of N=1 self-dual supergravity theories in (2,2) dimensions. One of them, which we call type I, can be obtained by chiral truncation of the simple supergravity in (2,2) dimensions. This truncation is very much like the one discussed above in the case of super Yang-Mills theory [7]. The remaining fields are $(e_\mu^a, \psi_{\mu-})$, the vierbein and the gravitino fields, respectively. The resulting field equations are

$$\begin{aligned} R_{cd}{}^{ab}(\omega(e)) - \tfrac{1}{2}\epsilon_{cdef}R^{efab}(\omega(e)) &= 0, \\ \psi_-^{ab} - \tfrac{1}{2}\epsilon^{abcd}\psi_{cd-} &= 0, \end{aligned} \tag{15}$$

where $R_{cd}{}^{ab}$ is the Riemann curvature and $\psi_{ab} = e_a^\mu e_b^\nu\,[D_\mu(\omega(e)\psi_\nu - D_\nu(\omega(e)\psi_\mu]$. The torsion-free spin connection $\omega(e)$ is the solution of the equation $D_\mu(\omega(e))e_\nu^a \equiv \partial_\mu e_\nu^a +$

$\omega_\mu{}^{ab} e_{\nu b} - \Gamma^\lambda_{\mu\nu}(g) e^a_\lambda = 0$, where $\Gamma^\lambda_{\mu\nu}(g)$ is the usual Christophel symbol. The solution is given by $\omega_{\mu ab} = \frac{1}{2} e^\nu_a (\partial_\mu e_{\nu b} - \partial_\nu e_{\mu b}) + \frac{1}{2} e_{\mu c} e^\rho_a e^\sigma_b \partial_\sigma e^c_\rho - a \leftrightarrow b$.

The type I self-dual supergravity system described in (15) has the local supersymmetry

$$\begin{aligned} \delta e_\mu{}^a &= \tfrac{1}{2} \bar{\epsilon}_+ \gamma^a \psi_{\mu -} \ , \\ \delta \psi_{\mu -} &= \mathcal{D}_\mu(\omega(e)) \epsilon_- \ . \end{aligned} \tag{16}$$

Closure of the algebra requires that the supersymmetry parameter ϵ_+ be covariantly constant, i.e. $D_\mu(\omega)\epsilon_+ = 0$. The integrability condition of this equation leads to the self-duality condition (15) for the Riemann curvature. For later convenience we record here the algebra of the transformations (16):

$$[\delta_{\epsilon_1}, \delta_{\epsilon_2}] = \delta_\xi + \delta_\Lambda + \delta_{\epsilon_3} \ , \tag{17}$$

where the composite general coordinate , Lorenz and supersymmetry parameters are given by

$$\begin{aligned} \xi^\mu &= \bar{\epsilon}_{2+} \gamma^\mu \epsilon_{1-} - 1 \leftrightarrow 2 \ , \\ \Lambda &= \xi^\nu \omega_{\nu ab} \ , \\ \epsilon_3 &= -\xi^\nu \psi_{\nu -} \ . \end{aligned} \tag{18}$$

In deriving this result, we have used the following formula for the supersymmetric variation of the spin connection:

$$\begin{aligned} \delta\omega_{\mu ab}(e) = & - \tfrac{1}{2} \bar{\epsilon}_+ \gamma_\mu \psi_{ab-} + \tfrac{1}{2} \bar{\epsilon}_+ \gamma_b \psi_{\mu a -} - \tfrac{1}{2} \bar{\epsilon}_+ \gamma_a \psi_{\mu b -} \\ & \left(\tfrac{1}{2} \bar{\psi}_{\mu -} \gamma_b D_a \epsilon_+ - \tfrac{1}{2} \bar{\psi}_{a-} \gamma_b D_\mu \epsilon_+ + \tfrac{1}{2} \bar{\psi}_{b-} \gamma_\mu D_a \epsilon_+ - a \leftrightarrow b \right) \ . \end{aligned} \tag{19}$$

and the Fierz rearrangement formulae: $\chi\bar{\psi} = -\frac{1}{2}\bar{\psi}\chi + \frac{1}{8}\bar{\psi}\gamma^{\mu\nu}\chi\gamma_{\mu\nu}$ for same chirality anti-commuting Majorana-Weyl spinors and $\chi\bar{\psi} = -\frac{1}{2}\bar{\psi}\gamma^\mu\chi\gamma_\mu$ for the case of opposite chiralities. A useful identity is: $\gamma^{\mu\nu} = -\frac{1}{2}\epsilon^{\mu\nu\rho\sigma}\gamma_{\rho\sigma}\gamma_5$.

Going back to the self-duality equations (15), we note that there are no fermionic corrections to the one involving the Riemann tensor. Nonetheless one verifies that these equations do transform into each other under the supersymmetry transformations (16), thanks to the gravitino equation $\gamma^{cab}\psi_{ab-} = 0$ and its consequences $\gamma^{ab}\psi_{ab-} = 0$ and $\gamma^a\psi_{ab-} = 0$. The gravitino equation has to be satisfied by closure requirements. It can also seen as a consequence of the self-duality equations (15) taken together with the Bianchi identities for the curvatures. This is similar to the fact that the self-duality condition on the Yang-Mills field strength plus the Bianchi identity imply the ordinary Yang-Mills field equation. Notice also that one can derive the self-duality equation from the gravitino equation. To see this we multiply $\gamma^{cab}\psi_{ab-} = 0$ by γ^d and use the equations $\gamma^{ab}\psi_{ab-} = 0$ and $\gamma^a\psi_{ab-} = 0$.

Some other aspects of the self-duality condition (15) are also noteworthy. From (15) and the identity $R_{[abc]d}(\omega(e)) = 0$, it follows that the Ricci-tensor vanishes $R_{ab}(\omega(e)) = 0$. We also know that Ricci-flat self-dual metrics are Kahler. Therefore, it follows that self-dual metrics are Kahler. For such metrics, using complex coordinates labeled by $i = 1, 2$, one can express the Ricci tensor as $R_{i\bar{j}} = \partial_i \partial_{\bar{j}} \ln \det g_{i\bar{j}}$, where $g_{i\bar{j}}$ is the metric. In terms of a Kahler potential K, the metric can be written as $g_{i\bar{j}} = \partial_i \partial_{\bar{j}} K$. Therefore, the vanishing of the Ricci tensor means that det $\partial_i \partial_{\bar{j}} K = 1$. This can be written as a Poisson bracket as $\epsilon^{ij}\{\partial_i K, \partial_j K\} = 2$, where the Poisson brackets of A and B is defined as $\{A, B\} = \epsilon^{\bar{i}\bar{j}} \partial_{\bar{i}} A \partial_{\bar{j}} B$. This equation is clearly invariant under symplectic diffeomorphisms of the form $\delta \partial_i K = \{\partial_i K, \Lambda\}$ where Λ is an arbitrary function. Note that the metric is not invariant under this transformation, and consequently, from a given metric which satisfies the Ricci flatness condition, one obtains a one parameter family of metrics which also obey this condition.

7. We now describe another type of N=1 self-dual supergravity theory, which we shall call type II, which can be obtained from the so-called new minimal formulation [9] of the off-shell supergravity (adapted to (2,2) signature). The new minimal multiplet consists of the vierbein ${e_\mu}^a$, the gravitini $\psi_{\mu\pm}$, the antisymmetric tensor field $B_{\mu\nu}$ and an SO(1,1) gauge field V_μ. It is to define the combinations

$$\begin{aligned} {\Omega_{\mu\pm}}^{ab} &= {\omega_\mu}^{ab}(e,\psi) \pm {\hat{H}_\mu}{}^{ab} \ , \\ V_{\mu+} &= V_\mu + \tfrac{1}{6} {\epsilon_\mu}^{abc} \hat{H}_{abc} \ , \end{aligned} \tag{20}$$

where the supercovariant spin-connection ${\omega_\mu}^{ab}(e,\psi)$ is defined as

$${\omega_\mu}^{ab}(e,\psi) = {\omega_\mu}^{ab}(e) - \tfrac{1}{2} \bar{\psi}_\mu \gamma^{[a} \psi^{b]} - \tfrac{1}{4} \bar{\psi}^a \gamma_\mu \psi^b \ , \tag{21}$$

with $\psi_\mu = \psi_{\mu+} + \psi_{\mu-}$, and the supercovariant curvatures are

$$\begin{aligned} \psi_{\mu\nu\pm} &= \mathcal{D}_\mu(\Omega_+, V_+)\psi_{\nu\pm} - \mathcal{D}_\nu(\Omega_+, V_+)\psi_{\mu\pm} \ , \\ \hat{H}_{\mu\nu\rho} &= \partial_{[\mu} B_{\nu\rho]} - \tfrac{3}{2} \bar{\psi}_{[\mu+} \gamma_\nu \psi_{\rho]-} \ , \\ \hat{F}_{ab}(V_+) &= 2\partial_{[a} V_{b]+} - \tfrac{1}{2} \bar{\psi}_{[a+} \gamma^\lambda \psi_{b]\lambda-} + \tfrac{1}{2} \bar{\psi}_{[a-} \gamma^\lambda \psi_{b]\lambda+} \ , \\ {\hat{R}_{cd}}{}^{ab}(\Omega_-) &= {R_{cd}}^{ab}(\Omega_-) - \bar{\psi}_{[c+} \gamma_{d]} \psi_-^{ab} - \bar{\psi}_{[c-} \gamma_{d]} \psi_+^{ab} \ . \end{aligned} \tag{22}$$

The derivative $\mathcal{D}_\mu$ on the supersymmetry parameters $\epsilon_\pm$ is given by

$$\mathcal{D}_\mu(\Omega_+, V_+)\epsilon_\pm = \Big(\partial_\mu - \tfrac{1}{4} {\Omega_{\mu+}}^{ab} \gamma_{ab} \mp V_{\mu+} \Big) \epsilon_\pm \tag{23}$$

The N=1 type II self-dual supergravity system consists of the following equations [5]

$$\begin{aligned} \psi_\pm^{ab} &= \tfrac{1}{2} \epsilon^{abcd} \psi_{cd\pm} \ , \\ {\hat{R}_{cd}}{}^{ab}(\Omega_-) &= \tfrac{1}{2} \epsilon^{abef} \hat{R}_{cdef}(\Omega_-) \ , \\ \hat{F}^{ab}(V_+) &= \tfrac{1}{2} \epsilon^{abcd} \hat{F}_{cd}(V_+) \ . \end{aligned} \tag{24}$$

These equations transform into each other under the following local supersymmetry transformations

$$\begin{aligned}\delta e_\mu{}^a &= \tfrac{1}{2}\bar\epsilon_+\gamma^a\psi_{\mu-} + \tfrac{1}{2}\bar\epsilon_-\gamma^a\psi_{\mu+}\\ \delta\psi_{\mu\pm} &= (\partial_\mu - \tfrac{1}{4}\Omega_{\mu+}{}^{ab}\gamma_{ab} \mp V_{\mu+})\epsilon_\pm\\ \delta B_{\mu\nu} &= \tfrac{3}{2}\bar\epsilon_+\gamma_{[\mu}\psi_{\nu]-} + \tfrac{3}{2}\bar\epsilon_-\gamma_{[\mu}\psi_{\nu]+}\\ \delta V_\mu &= \tfrac{1}{8}\bar\epsilon_+\gamma_\mu\gamma^{ab}\psi_{ab-}\end{aligned}\tag{25}$$

It is interesting to observe [5] that the last self-duality condition in (24) together with the Bianchi identity $D_{[a}\hat F_{bc]}(V_+) + \cdots = 0$ imply the field equation $D_a\hat F^{ab}(V_+) + \cdots = 0$, which we recognize as one of the field equations that follow from the (2,2) version of the R^2-type action constructed in [10] given by

$$\begin{aligned}e^{-1}\mathcal{L}(R^2) =& \tfrac{1}{4}R_{\mu\nu}{}^{ab}(\Omega_-)R^{\mu\nu ab}(\Omega_-) - 2\hat F_{ab}(V_+)\hat F^{ab}(V_+) + \tfrac{1}{2}\bar\psi^{ab}\gamma^\mu D_\mu(\Omega_-,V)\psi_{ab}\\ &+ \tfrac{1}{8}\bar\psi_\mu\gamma^{cd}\gamma^\mu\psi_{ab}\big\{R_{cd}{}^{ab}(\Omega_-) + \hat R_{cd}{}^{ab}(\Omega_-)\big\}\end{aligned}\tag{26}$$

where $\psi_\mu = \psi_{\mu+} + \psi_{\mu-}$ and the Ω_- covariantization in the kinetic term of the gravitino curvature acts both on the spinor as well as the vector indices of ψ^{ab}. By supersymmetry it then follows that the field equations for the remaining fields $e_\mu{}^a, \psi_{\mu\pm}$ and $B_{\mu\nu}$ must also be satisfied.

The N=2 and N=4 self-dual supergravity theories in (2,2) dimensions can also be obtained from their non-self-dual counterparts (which in turn can be obtained by a Wick rotation of their (3,1) versions) by chirality/self-duality restrictions [8,2]. The case of N=8 supergravity, just as in the case of N=4 super Yang-Mills, turns out to be more subtle, and has been treated in [2].

8. In [2], it was found that a self-dual version of *gauged* N=8 supergravity arsies as an effective theory of closed N=2 superstrings. The relevant global superalgebra, in the case of non-self dual extended supergravity in (2,2) dimensions, is $OSp(N|4)$, whose bosonic subalgebra is $SO(3,2)\oplus SO(n,N-n)$. It can be obtained from the superconformal algebra $SL(N|4)$, which has the bosonic subalgebra $SO(3,3)\oplus SO(1,1)\oplus SL(N,R)$, by a suitable restriction. In the resulting algebra, the SO(3,2) generators can be decomposed with respect to the SO(2,2) subalgebra as $(M_{\mu\nu}, P_\mu)$ and in addition to the obvious commutation rules one has

$$\begin{aligned}&\{Q^i_+, Q^j_+\} = \gamma^{\mu\nu}CM^-_{\mu\nu}\eta^{ij} + \tfrac{1}{2}(1+\gamma_5)CT^{ij}\ ,\\ &\{Q^i_+, Q^j_-\} = \tfrac{1}{2}(1+\gamma_5)\gamma^\mu CP_\mu\eta^{ij}\ ,\\ &\{Q^i_-, Q^j_-\} = \gamma^{\mu\nu}CM^+_{\mu\nu}\eta^{ij} + \tfrac{1}{2}(1-\gamma_5)CT^{ij}\ ,\\ &\quad[P_\mu, Q^i_\pm] = \tfrac{1}{2}\gamma_\mu Q^i_\mp\ , \qquad\qquad [M^\mp_{\mu\nu}, Q_\pm] = \tfrac{1}{2}\gamma_{\mu\nu}Q_\pm\ ,\end{aligned}\tag{27}$$

where η^{ij} is the invariant tensor, and T^{ij} are the generators, of $SO(n,N-n)$, and $M^\pm_{\mu\nu}$ are the self-dual and anti-self dual pieces of the Lorentz generators.

The gauged N=8 supergravity in (2,2) dimension is rather complicated. Its self-dual restriction is in some ways similar to the case of N=4 super Yang-Mills theory, where propagating Lagrange multipliers are needed to impose the self-duality condition. An important property of gauged supergravities is that they contain a cosmological constant. While the cosmological constant is necessary in gauged supergravity theories, gauging is not necessary in order to introduce a cosmological constant. The simplest example in four dimensions is N=1 supergravity with cosmological constant [11]. It is instructive to consider the consequences of its Wick rotation to (2,2) dimensions and its chiral truncation. The transformation rules in (2,2) dimensions are

$$\begin{aligned} \delta e_\mu^a &= \bar\epsilon_+\gamma^a\psi_{\mu-} + \bar\epsilon_-\gamma^a\psi_{\mu+}\ , \\ \delta\psi_{\mu+} &= D_\mu(\hat\omega)\epsilon_+ + m\gamma_\mu\epsilon_-\ , \\ \delta\psi_{\mu-} &= D_\mu(\hat\omega)\epsilon_- + m\gamma_\mu\epsilon_+\ , \end{aligned} \tag{28}$$

where m is an arbitrary mass parameter, and $\hat\omega_{\mu ab}$ equals $\omega(e)_{\mu ab}$ plus fermionic torsion which consists of bilinears in the gravitino field. The closure of this algebra requires the field equations that follow from the supergravity action plus a gravitino mass term and a cosmological constant term. In particular, $\gamma^{\mu\nu\rho}\psi_{\nu\rho-} \sim m\gamma^{\mu\nu}\psi_{\nu+}$ and $\gamma^{\mu\nu\rho}\psi_{\nu\rho+} \sim m\gamma^{\mu\nu}\psi_{\nu-}$.

Now let us consider the chiral truncation $\psi_+ = 0$. From (28) we get the condition $D_\mu(\omega(e))\epsilon_+ + m\gamma_\mu\epsilon_- = 0$, which determines ϵ_- in terms of ϵ_+. However, we run into a problem with the gravitino field equation: $\gamma^{\mu\nu}\psi_{\nu-} = 0$, and hence $\psi_{\mu-} = 0$. Interestingly enough, there is another route one can take, which turns out to imply that the torsion-free self-dual N=1 supergravity equations actually admit a one parameter extension of the super Poincaré algebra as a symmetry algebra. To see this, let us consider the folowing transformations

$$\begin{aligned} \delta e_\mu^a &= \bar\epsilon_+\gamma^a\psi_{\mu-}\ , \\ \delta\psi_{\mu-} &= D_\mu(\omega(e))\epsilon_- + m\gamma_\mu\epsilon_+\ . \end{aligned} \tag{29}$$

Remarkably, these transformation rules do form a closed algebra provided that $D_\mu(\omega(e))\epsilon_+ = 0$ and the self-dual supergravity equations (15) are satisfied. The algebra of these transformation is

$$[\delta_{\epsilon_1},\delta_{\epsilon_2}] = \delta_\xi + \delta_{\Lambda'} + \delta_{\epsilon_3}\ , \tag{30}$$

where the composite general coordinate, Lorenz and supersymmetry parameters are given by

$$\begin{aligned} \xi^\mu &= \bar\epsilon_{2+}\gamma^\mu\epsilon_{1-} - 1 \leftrightarrow 2\ , \\ \Lambda' &= \xi^\nu\omega_{\nu ab} + 2m\bar\epsilon_{2+}\gamma_{ab}\epsilon_{1+}\ , \\ \epsilon_3 &= -\xi^\nu\psi_{\nu-}\ . \end{aligned} \tag{31}$$

One can check that $\delta_{\epsilon_3}e_\mu^a = 0$ and that $\delta_{\Lambda'}\psi_{\mu-} = \delta_\Lambda\psi_{\mu-}$. We see that unlike in the case of the algebra (17), here even in the global limit we still have an (anti) self-dual Lorentz rotation. Noting that $[\delta_\epsilon,\delta_\xi] = \delta_{\epsilon'}$ and $[\delta_\epsilon,\delta_\Lambda] = \delta_{\epsilon''}$ where $\epsilon' = -\xi^\mu\partial_\mu\epsilon$ and $\epsilon'' = -\frac{1}{4}\Lambda_{ab}\gamma^{ab}\epsilon$, we see that the global limit of the full algebra contains the generators $(Q_\pm, M^-_{\mu\nu}, P_\mu)$, obeying the algebra $\{Q_+, Q_+\} = \gamma^{\mu\nu} C M^-_{\mu\nu}$, $\{Q_+, Q_-\} = \frac{1}{2}(1+\gamma_5)\gamma^\mu C P_\mu$, and $[M^-_{\mu\nu}, Q_+] = \frac{1}{2}\gamma_{\mu\nu} Q_+$, plus the commutators of the ordinary Poincaré algeba, with all the other (anti) commutators vanishing.

Acknowledments

I thank Mike Duff, Chris Pope and Larry Romans for stimulating discussions.

References

1. H. Ooguri and C. Vafa, Mod. Phys. Lett. **A5** (1990) 1389; Nucl. Phys. **B361** (1991) 469; Nucl. Phys. **B367** (1991) 83.
2. W. Siegel, preprints, ITP-SB-92-24, 31,53.
3. See, for example, the review article by:

   R.S. Ward in *Field Theory, Quantum Gravity and Strings*, Vol. 2, p. 106, eds. H.J. De Vega and N. Sanchez.
4. S.J. Gates Jr., H. Nishino and S.V. Ketov, preprint, UMDEPP 92-51.
5. E. Bergshoeff and E. Sezgin, preprint, UG-4/92, CTP TAMU-46/92.
6. For the properties of spinors in arbitrary dimensions, see:

   T. Kugo and P.K. Townsend, *Nucl. Phys.* **B221** (1983) 357.
7. S.M. Christensen and M.J. Duff, Nucl. Phys. **B159**(1979) 429;

   S.M. Christensen, S. Deser, M.J. Duff and M.T. Grisaru, Phys. Lett. **84B** (1979) 411; R.E. Kallosh, JETP Lett. **30** (1979)172,449; Nucl. Phys. **B165** (1980) 119.
8. S.J. Gates Jr., H. Nishino and S.V. Ketov, preprints, UMDEPP 92-163,171,187,211.
9. M.F. Sohnius and P.C. West, Phys. Lett. **105B** (1981) 353.
10. M. de Roo, A. Wiedemann and E. Zijlstra, Class. Quantum Grav. **7** (1990) 1181.
11. P.K. Townsend, Phys. Rev. **D15** (1977) 2808.